

Effect of developmental pesticide exposure in Prairie voles as a model of neurodevelopmental disorders

Nilanjana Saferin^{1*}, Sangeetha Pillai¹, Dena Hasan¹, Isa Malik¹, Ibrahim Haseeb¹, Asha Neifer², Kari Neifer², James Burkett, PhD³

¹College of Medicine and Life Sciences, The University of Toledo, Toledo, Ohio 43614

²Department of Physiology and Pharmacology, The University of Toledo, Toledo, Ohio 43614

³Department of Neurosciences, The University of Toledo, Toledo, Ohio 43614

*Corresponding author: nilanjana.chakrabarti@utoledo.edu

Keywords: Pyrethroids, Deltamethrin, Prairie Voles, Neurodevelopmental Disorders

Published: 22 May 2024

Background: Neurodevelopmental disorders (NDDs) are a class of lifelong incurable disorders with few treatments are biomarkers. They share common comorbidities including deficits in communication and learning, repetitive behavior, and hyperactivity such as attention deficit hyperactivity disorder (ADHD), Autism, and Developmental disability. The incidence of NDDs has been rapidly rising affecting almost 17% of children in the US. Heritability is still a crucial factor in the etiology of NDDs, however large meta-analysis data now consider environmental impact as a major contributor. Recent epidemiological studies have shown the effect of pyrethroid pesticides on pregnant women and risk factors associated with the proper brain development in the children. Pyrethroids are common household insecticides widely used in the US and considered relatively “safe” by the Environmental Protection Agency (EPA). Despite growing evidence of the complex gene-environment interaction in the etiology of NDDs, very few environmental factors have been studied.

Methods: Previously, our lab has studied the effects of deltamethrin (pyrethroid) in mice. Metabolomic analysis on whole male brain samples provided suggested a disruption in folate metabolism pathway as a result of developmental pesticide exposure. Our research aims to look at the effect of developmental pyrethroid exposure (DPE) in prairie voles as a model of neurodevelopmental disorders, and folic acid supplementation as a potential therapeutic strategy.

Results: We exposed pregnant vole dams to 3mg/kg of deltamethrin two weeks prior to pregnancy, during pregnancy and throughout lactation. A subset of pesticide exposed voles were supplemented with 5-MTHF (folate vitamer). Following weaning, the offspring grow to adulthood and are subjected to a battery of behavioral tests to look at deficits in five different domains namely communication, cognition, social interaction, repetitive behavior, and locomotion. After concluding behavioral tests their brains are harvested along with other tissues for further studies. In order to understand molecular disruptions in the folate metabolic enzymatic pathway, we conducted western blot analysis in five participating enzymes in the folate metabolic pathway, namely Dihydrofolate Reductase (DHFR), Methyltetrahydrofolate

Reductase (MTHFR), Serine hydroxymethyl transferase (SHMT), Methyl tetrahydrofolate dehydrogenase (MTHFD), Methionine synthase (MTRR), and Folate receptor alpha (FOLR1).

Conclusions: Our results show that developmentally exposed prairie voles had deficits in communication, cognition, repetitive behavior, and locomotion (hyperactivity). Previously harvested brains from the mice study were used to look at disruptions in molecular mechanisms because of DPE. Our data also suggest that folate supplementation alleviates some of the behavioral deficits. Disruptions in enzymatic expression was observed in FOLR1, MTHFR, and SHMT1. Such disruptions could be an effect of treatment or a compensatory mechanism for changes caused by developmental pesticide exposure.